

DIVERTICULOSIS & DIVERTICULITIS




What is diverticulosis and diverticulitis?

Many people have small pouches in the lining of the colon, or large intestine, which bulge outward through weak spots. Each pouch is called a *diverticulum*. Multiple pouches are called *diverticula*. The condition of having diverticula is called *diverticulosis*. About 10 percent of Americans older than 40 have diverticulosis.¹ The condition becomes more common as people age. About half of all people older than 60 have diverticulosis.²

Diverticula are most common in the lower portion of the large intestine, called the sigmoid colon. When the pouches become inflamed, the condition is called *diverticulitis*. Ten to 25 percent of people with diverticulosis get diverticulitis.³ Diverticulosis and diverticulitis together are called *diverticular disease*.

¹Bogardus ST. What do we know about diverticular disease? A brief overview. *Journal of Clinical Gastroenterology*. 2006;40:S108–S111.
²Ibid.
³Ibid.



What are the symptoms of diverticulosis and diverticulitis?

Diverticulosis—Most people with diverticulosis do not have any discomfort or symptoms. However, some people may experience crampy pain or discomfort in the lower abdomen, bloating, and constipation. Other conditions such as irritable bowel syndrome and stomach ulcers cause similar problems, so the symptoms do not always mean a person has diverticulosis. People with chronic symptoms should visit their doctor or health care provider.

Diverticulitis—The most common symptom of diverticulitis is abdominal pain. The most common sign on examination is tenderness in the lower left side of the abdomen. Usually, the pain is severe and comes on suddenly, but it can also be mild and become worse over several days. The intensity of the pain can fluctuate. A person may experience cramping, nausea, vomiting, fever, chills, or a change in bowel habits.

What are the complications of diverticulitis?

Diverticulitis can lead to bleeding; infections; small tears, called perforations; or blockages in the colon. These complications always require treatment to prevent them from progressing and causing serious illness.

Bleeding—Rectal bleeding from diverticula is a rare complication. Doctors believe the bleeding is caused by a small blood vessel in a diverticulum that weakens and then bursts. When diverticula bleed, blood may appear in the toilet or in the stool. Bleeding can be severe, but it may stop by itself and not require treatment. A person who has bleeding from the rectum—even a small amount—should see a doctor right away. Often, colonoscopy is used to identify the site of bleeding and stop the bleeding. Sometimes the doctor injects dye into an artery—a procedure called angiography—to identify and treat diverticular bleeding. If the bleeding does not stop, surgery may be necessary to remove the involved portion of the colon.

Abscess, Perforation, and Peritonitis-Diverticulitis may lead to infection, which often clears up after a few days of treatment with antibiotics. If the infection gets worse, an abscess may form in the wall of the colon.

An abscess is a localized collection of pus that may cause swelling and destroy tissue. If the abscess is small and remains in the wall of the colon, it may clear up after treatment with antibiotics. If the abscess does not clear up with antibiotics, the doctor may need to drain it using a catheter—a small tube—placed into the abscess through the skin. After giving the patient numbing medicine, the doctor inserts the needle through the skin until reaching the abscess and then drains the fluid through the catheter. This process may be guided by sonography or x ray.

Infected diverticula may develop perforations. Sometimes the perforations leak pus out of the colon and form a large abscess in the abdominal cavity, a condition called peritonitis. A person with peritonitis may be extremely ill with nausea, vomiting, fever, and severe abdominal tenderness. The condition requires immediate surgery to clean the abdominal cavity and remove the damaged part of the colon. Without prompt treatment, peritonitis can be fatal.

Fistula-A fistula is an abnormal connection of tissue between two organs or between an organ and the skin. When damaged tissues come into contact with each other during infection, they sometimes stick together. If they heal that way, a fistula may form. When diverticulitis-related infection spreads outside the colon, the colon's tissue may stick to nearby tissues. The organs usually involved are the bladder, small intestine, and skin.

The most common type of fistula occurs between the bladder and the colon. This type of fistula affects men more often than women. It can result in a severe, long-lasting infection of the urinary tract. The problem can be corrected with surgery to remove the fistula and the affected part of the colon.

Intestinal Obstruction-Scarring caused by infection may lead to partial or total blockage of the intestine, called intestinal obstruction. When the intestine is blocked, the colon is unable to move bowel contents normally. If the intestine is completely blocked, emergency surgery is necessary. Partial blockage is not an emergency, so the surgery to correct it can be planned.

What causes diverticular disease?

Although not proven, the dominant theory is that a low-fiber diet causes diverticular disease. The disease was first noticed in the United States in the early 1900s, around the time processed foods were introduced into the American diet. Consumption of processed foods greatly reduced Americans' fiber intake.

Diverticular disease is common in developed or industrialized countries—particularly the United States, England, and Australia—where low-fiber diets are consumed. The disease is rare in Asia and Africa, where most people eat high-fiber diets.

Fiber is the part of fruits, vegetables, and grains that the body cannot digest. Some fiber, called soluble fiber, dissolves easily in water. It takes on a soft, jelly-like texture in the intestines. Insoluble fiber passes almost unchanged through the intestines. Both kinds of fiber help prevent constipation by making stools soft and easy to pass.

Constipation—or hard stool—may cause people to strain when passing stool during a bowel movement. Straining may cause increased pressure in the colon, which may cause the colon lining to bulge out through weak spots in the colon wall. These bulges are diverticula.

Lack of exercise also may be associated with a greater risk of forming diverticula, although the reasons for this are not well understood.

Doctors are not certain what causes diverticula to become inflamed. The inflammation may begin when bacteria or stool are caught in the diverticula. An attack of diverticulitis can develop suddenly and without warning.

How is diverticular disease diagnosed?

To diagnose diverticular disease, the doctor asks about medical history, does a physical exam, and may perform one or more diagnostic tests. Because most people do not have symptoms, diverticulosis is often found through tests ordered for another ailment. For example, diverticulosis is often found during a colonoscopy done to screen for cancer or polyps or to evaluate complaints of pain or rectal bleeding.

When taking a medical history, the doctor may ask about bowel habits, pain, other symptoms, diet, and medications. The physical exam usually involves a digital rectal exam. To perform this test, the doctor inserts a gloved, lubricated finger into the rectum to detect tenderness, blockage, or blood. The doctor may check stool for signs of bleeding and test blood for signs of infection. If diverticulitis is suspected, the doctor may order one of the following radiologic tests:

- **Abdominal ultrasound.** Sound waves are sent toward the colon through a handheld device that a technician glides over the abdomen. The sound waves bounce off the colon and other organs, and their echoes make electrical impulses that create a picture—called a sonogram—on a video monitor. If the diverticula are inflamed, the sound waves will also bounce off of them, showing their location.
- **Computerized tomography (CT) scan.** The CT scan is a noninvasive x ray that produces cross-section images of the body. The doctor may inject dye into a vein and the person may be given a similar mixture to swallow. The person lies on a table that slides into a donut-shaped machine. The dye helps to show complications of diverticulitis such as perforations and abscesses.

How is diverticular disease treated?

A high-fiber diet and pain medications help relieve symptoms in most cases of diverticulosis. Uncomplicated diverticulitis with mild symptoms usually requires the person to rest, take oral antibiotics, and be on a liquid diet for a period of time. Sometimes an attack of diverticulitis is serious enough to require a hospital stay, intravenous (IV) antibiotics, and possibly surgery.

Diverticulosis-Increasing the amount of fiber in the diet may reduce symptoms of diverticulosis and prevent complications such as diverticulitis. Fiber keeps stool soft and lowers pressure inside the colon so that bowel contents can move through easily. The American Dietetic Association recommends consuming 20 to 35 grams of fiber each day. The table “What foods have fiber?” shows the amount of fiber in some foods that a person can easily add to the diet.

The doctor may also recommend taking a fiber product such as methylcellulose (Citrucel) or psyllium (Metamucil) one to three times a day. These products are available in powder, pills, or wafers, and provide 2 to 3.5 grams of fiber per dose. Fiber products should be taken with at least 8 ounces of water.

Avoidance of nuts, popcorn, and sunflower, pumpkin, caraway, and sesame seeds has been recommended by physicians out of fear that food particles could enter, block, or irritate the diverticula. However, no scientific data support this treatment measure. Eating a high-fiber diet is the only requirement highly emphasized across the medical literature. Eliminating specific foods is not necessary. The seeds in tomatoes, zucchini, cucumbers, strawberries, and raspberries, as well as poppy seeds, are generally considered harmless. People differ in the amounts and types of foods they can eat. Decisions about diet should be made based on what works best for each person. Keeping a food diary may help identify what foods may cause symptoms.

If cramps, bloating, and constipation are problems, the doctor may prescribe a short course of pain medication. However, some pain medications actually cause constipation.

Diverticulitis-Treatment for diverticulitis focuses on clearing up the inflammation and infection, resting the colon, and preventing or minimizing complications.

Depending on the severity of symptoms, the doctor may recommend bed rest, oral antibiotics, a pain reliever, and a liquid diet. If symptoms ease after a few days, the doctor will recommend gradually increasing the amount of high-fiber foods in the diet.

Severe cases of diverticulitis with acute pain and complications will likely require a hospital stay. Most cases of severe diverticulitis are treated with IV antibiotics and a few days without food or drink to help the colon rest. In some cases, surgery may be necessary.

What foods have fiber?

Examples of foods that have fiber include

Breads, cereals, and beans

- 1/2 cup of navy beans= 9.5 grams
- 1/2 cup of kidney beans= 8.2 grams
- 1/2 cup of black beans=7.5 grams

Whole-grain cereal, cold

- 1/2 cup of All-Bran= 9.6 grams
- 3/4 cup of Total= 2.4 grams
- 3/4 cup of Post Bran Flakes= 5.3 grams


1 packet of whole-grain cereal, hot= 3.0 grams
(oatmeal, Wheatena)

1 whole-wheat English muffin4.4 grams



Fruits

- 1 medium apple, with skin3.3 grams
- 1 medium pear, with skin4.3 grams
- 1/2 cup of raspberries4.0 grams
- 1/2 cup of stewed prunes3.8 grams



Vegetables

- 1/2 cup of winter squash2.9 grams
- 1 medium sweet potato with skin4.8 grams
- 1/2 cup of green peas4.4 grams
- 1 medium potato with skin3.8 grams
- 1/2 cup of mixed vegetables4.0 grams

1 cup of cauliflower 2.5 grams

1/2 cup of spinach 3.5 grams

1/2 cup of turnip greens 2.5 grams

Source: U.S. Department of Agriculture and U.S. Department of Health and Human Services, *Dietary Guidelines for Americans*, 2005

When is surgery necessary for diverticulitis?

If symptoms of diverticulitis are frequent, or the patient does not respond to antibiotics and resting the colon, the doctor may advise surgery. The surgeon removes the affected part of the colon and joins the remaining sections. This type of surgery—called colon resection—aims to prevent complications and future diverticulitis. The doctor may also recommend surgery for complications such as a fistula or partial intestinal obstruction.

Immediate surgery may be necessary when the patient has other complications, such as perforation, a large abscess, peritonitis, complete intestinal obstruction, or severe bleeding. In these cases, two surgeries may be needed because it is not safe to rejoin the colon right away. During the first surgery, the surgeon cleans the infected abdominal cavity, removes the portion of the affected colon, and performs a temporary colostomy, creating an opening, or stoma, in the abdomen. The end of the colon is connected to the opening to allow normal eating while healing occurs. Stool is collected in a pouch attached to the stoma. In the second surgery several months later, the surgeon rejoins the ends of the colon and closes the stoma.

Points to Remember

- Diverticulosis occurs when small pouches called diverticula bulge outward through weak spots in the colon, or large intestine.
- Most people with diverticulosis never have any discomfort or symptoms.
- Diverticula form when pressure builds inside the colon wall, usually because of constipation.
- The most likely cause of diverticulosis is a low-fiber diet because it increases constipation and pressure inside the colon.
- For most people with diverticulosis, eating a high-fiber diet is the only treatment needed.
- Fiber intake can be increased by eating whole-grain breads and cereals; fruits like apples and pears; vegetables like peas, spinach, and squash; and starchy vegetables like kidney and black beans.
- Diverticulitis occurs when the pouches become inflamed and cause pain and tenderness in the lower left side of the abdomen.
- Diverticulitis can lead to bleeding; infections; small tears, called perforations; or blockages in the colon. These complications always require treatment to prevent them from progressing and causing serious illness.
- Severe cases of diverticulitis with acute pain and complications will likely require a hospital stay. When a person has complications or does not respond to medication, surgery may be necessary.

Hope through Research

Coastal Digestive Care Center, LLC.
www.coastaldigestivecare.com
860-447-0290

The National Institute of Diabetes and Digestive and Kidney Diseases and the National Cancer Institute sponsor research programs to investigate diverticulosis and diverticulitis.

Investigation continues in several areas, including

- a possible link between diverticular disease and inflammatory bowel disease
- the management of recurrent diverticular disease
- the use of probiotics in the prevention and treatment of diverticular disease
- indications for surgery for uncomplicated diverticulitis

Participants in clinical trials can play a more active role in their own health care, gain access to new research treatments before they are widely available, and help others by contributing to medical research. For information about current studies, visit www.ClinicalTrials.gov.

The U.S. Government does not endorse or favor any specific commercial product or company. Trade, proprietary, or company names appearing in this document are used only because they are considered necessary in the context of the information provided. If a product is not mentioned, the omission does not mean or imply that the product is unsatisfactory.

For More Information:

American College of Gastroenterology

P.O. Box 342260
Bethesda, MD 20827-2260
Phone: 301-263-9000
Fax: 301-263-9025
Internet: www.acg.gi.org

American Gastroenterological Association

4930 Del Ray Avenue
Bethesda, MD 20814
Phone: 301-654-2055
Fax: 301-654-5920
Email: member@gastro.org
Internet: www.gastro.org

You may also find additional information about this topic by visiting MedlinePlus at www.medlineplus.gov.

American Society of Colon and Rectal Surgeons

85 West Algonquin Road, Suite 550
Arlington Heights, IL 60005
Phone: 847-290-9184
Fax: 847-290-9203
Email: ascrs@fascrs.org
Internet: www.fascrs.org

International Foundation for Functional Gastrointestinal Disorders

P.O. Box 170864
Milwaukee, WI 53217-8076
Phone: 1-888-964-2001 or 414-964-1799
Fax: 414-964-7176
Email: iffgd@iffgd.org
Internet: www.iffgd.org

The National Digestive Diseases Information Clearinghouse (NDDIC) is a service of the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). The NIDDK is part of the National Institutes of Health of the U.S. Department of Health and Human Services. Established in 1980, the Clearinghouse provides information about digestive diseases to people with digestive disorders and to their families, health care professionals, and the public. The NDDIC answers inquiries, develops and distributes publications, and works closely with professional and patient organizations and Government agencies to coordinate resources about digestive diseases.

Publications produced by the Clearinghouse are carefully reviewed by both NIDDK scientists and outside experts. This publication was reviewed by Lawrence R. Schiller, M.D., Baylor University Medical Center at Dallas.

This publication is not copyrighted. The Clearinghouse encourages users of this publication to duplicate and distribute as many copies as desired.

Coastal Digestive Care Center, LLC.
www.coastaldigestivecare.com
860-447-0290